

Spider mites

Two-spotted spider mite – *Tetranychus urticae*

Bean spider mite – *T. luedeni*

Strawberry spider mite – *T. lambi*

The two-spotted spider mite is the main pest species, the other two species rarely colonise cotton and seldom cause economic damage. Even in high numbers, *T. lambi* infestations still result in very low levels of damage.

Damage symptoms

Nymphs and adults cause mid to late season damage to leaves. Leaves firstly redden and with continued feeding will become necrotic.

Sampling

'Sampling protocols for mites in cotton' is presented in full on page 22.

Sample what?

Look for the presence of any mite stages. Eggs and immature stages are difficult to see with the naked eye, so a hand lens should be used. Mites infest the underside of leaves. Sample the oldest leaf when plants are very young. As plants grow, choose leaves that are from 3, 4 or 5 nodes below the plant terminal.

Check which species is present. Two-spotted spider mite is pale green and has 2 distinct dark green spots on either side. Adults of bean spider mite are a dark red colour. Strawberry spider mite is smaller than the other two spider mites. Their bodies are pale green with 3 dark green spots on either side. They cause very little damage.



Two spotted mite with egg (mite is 0.5 mm long).
(Lewis Wilson, CSIRO)

Frequency

Sample at least weekly. Begin at seedling emergence.

Sample more frequently if mite populations begin to increase, or if conditions are hot and dry, or if sprays which eliminate predators are used.

Methods

Presence/absence sampling allows many plants to be sampled quickly, thus increasing the likelihood of finding mites if they are present. It is helpful to plot the development of mite populations on a graph. This allows changes in mite population to be seen at a glance. The detailed sampling protocol for monitoring mite populations is on page 20.

Thresholds

A general threshold of 30% of plants infested is advocated through the bulk of the season (squaring to first open boll). Yield loss due to mites depends on when mite populations begin to increase and how quickly they increase.

Seedling emergence to squaring

Mites are normally suppressed by predators, especially by thrips during this period. Mite populations only need to be controlled if they begin to increase, which indicates that natural controls are not keeping them in check. Use Table 4 on page 21 to determine whether the rate of increase warrants control.

Squaring to first open boll

Control if mite populations increase at greater than 1% of plants infested per day in two consecutive checks, or if more than 30% of plants are infested. Refer to Table 4 on page 21 for details.

First open bolls to 20% open bolls

Control is only warranted if mites are well established (greater than 60% plants infested) and are increasing rapidly (faster than 3% of plants infested per day). Refer to Table 4 on page 21 for details.

Crop exceeds 20% open bolls

Control is no longer warranted.

Mite Yield Loss estimator on the web

A simple relationship has been developed which allows prediction of yield loss from mites based on knowledge of the rate of increase in the population and the time remaining until defoliation. Record keeping and calculating can be simplified by using the Mite Yield Loss Estimator in CottASSIST on the web. Examples of charts generated by this tool are presented on page 22:

Mite population %. This is the percentage of leaves infested with mites.

Average rate of change. This is an average of the rates of change recorded for successive mite samples. Compared with the rate of change that you would expect if the yield loss from the mite population was 4%. This value (4%) is roughly when yield loss from mites would justify control, based on loss of revenue and cost of control. This may need to be adjusted for your particular situation.

Yield loss %. The yield loss calculation is based on the current percentage of plants infested with mites, the rate of change of the mite population and the number of days remaining in the season depending on the region. In general, zero or negative change in mite populations indicates that something has adversely affected population development such as mite spray, beneficials eating mites, heavy rainfall or a combination of these factors.

Mite yield reduction charts

As an alternative to the web tool, 'look-up' charts have been provided in Table 4, page 21 for areas with different season lengths:

Warmer – Bourke, Central Queensland, Macintyre Valley, St George and Walgett

Average – Dalby, Gwydir Valley, Lockyer Valley and Lower Namoi Valley

Cooler – Boggabri, Breeza, Cecil Plains – Pittsworth and Macquarie Valley

The charts use the rate of increase of the mite population. This is calculated by dividing the change in the percentage of plants infested between consecutive checks by the number of days between the checks. For example, if a field had 10% of plants infested a week ago and 24% infested now, this gives a rate of increase of 2% of plants infested per day.

To use the charts

1. Select the chart appropriate for your region.
2. Go to the section that is closest to the current infestation level of the field i.e. 10%, 30% or 60%.
3. Go to the column with the rate of increase closest to that of the mite population in the field.
4. Look down this column to the value that corresponds with the current age of the crop.

This value is the predicted yield loss that the mite population is likely to cause if left uncontrolled. It must be stressed that these charts only provide a guide for potential yield losses caused by mites.

You will need to take into account the vigour of the crop, other pests (you may be about to spray with a pyrethroid which may flare mites) and the conditions (that is, mites are generally favoured by hot dry conditions). Differences between the more mite resistant 'okra' leaf varieties and the normal leaf varieties are built into the charts.

Key beneficial insects

Predators – thrips, minute two-spotted ladybird, mite-eating ladybird, damsel bug, big-eyed bug, brown lacewing adults, brown smudge bug, apple dimpling bug, tangleweb spiders.

Selecting a miticide

The miticide products registered for the control of spider mites in cotton in Australia are presented in Table 5 on page 22.

Amitraz and endosulfan are insecticides that may be used for the control of *Helicoverpa spp.* early in the cotton season. When used for this purpose, these products tend to slow, or suppress, the development of mite populations that may also be in the field. Conversely, mite infestations may increase after the application of some broad spectrum insecticides used for *Helicoverpa* or mirid control, such as synthetic pyrethroids, and organophosphates. Known as 'mite flaring', this occurs because these sprays kill key beneficial species allowing mite populations to flourish.

Survival strategies

Resistance profile – two-spotted spider mite

WIDESPREAD, HIGH LEVELS OF RESISTANCE	WIDESPREAD, LOW/MOD LEVELS OF RESISTANCE
bifenthrin (SP) profenofos (OP)	chlorfenapyr
OCCASIONAL DETECTION OF HIGH LEVELS OF RESISTANCE	OCCASIONAL DETECTION OF LOW LEVELS OF RESISTANCE
	propargite

Abamectin resistance has occasionally been detected at high levels in two-spotted spider mite in horticulture, but not in cotton. Populations of the bean spider mite and the strawberry spider mite have not been tested for resistance to miticides because they are so rarely a problem in cotton.

Preventing resistance in two-spotted mites is complicated by the fact that most chemicals used for their control in cotton are also registered for the control of other pests, such as aphids or *Helicoverpa*. The bifenthrin and chlorfenapyr resistance that has developed in mites in recent years has occurred largely due to the use of these compounds against other pests.

Overwintering habit

Mites mostly survive the winter in cotton growing areas as active colonies. While the lifecycle slows in cool temperatures, mites are adapted to exploit ephemeral hosts and to produce large numbers of offspring, especially as conditions warm up in spring.

Alternative hosts

Preferred winter weed hosts are turnip weed, marshmallow, deadnettle, medics, wireweed and sowthistle. Alternative winter and spring host crops include safflower, faba beans and field peas.

Further Information

CSIRO Plant Industries, Narrabri

Lewis Wilson: (02) 6799 1550 or 0427 991 550

I&I NSW, Camden

Grant Herron: (02) 4640 6471

SAMPLING PROTOCOLS FOR MITES IN COTTON

Population Monitoring

1. Walk into the field about 40 m. (Early in the season it is also advisable to sample near the field edges to see if significant influxes of mites have occurred).
2. Take a leaf from the first plant on the right or left. The leaf should be from the third, fourth or fifth main-stem node below the terminal. If the plant has less than three leaves, sample the oldest. Note that early in the season, up to the point that the plant has about five true leaves, it is simplest to pull out whole plants.
3. Walk five steps and take a leaf from the next plant, on the opposite side to the previous one, and so on until you have 50 leaves. (Wait until you have collected all the leaves before scoring them).
4. Once all the leaves have been collected score each leaf by turning it over, looking at the underside, firstly near the stalk, then scanning the rest of the leaf. If mites of any stage (eggs or motiles) are present score the leaf as infested. A hand lens will be needed to see mite eggs because they cannot be seen with the naked eye.
5. Repeat this simple procedure at several widely separated places in the field to allow for differences in mite abundance within the field. Depending on the size of the field, 4–6 sites are needed to obtain a good estimate of mite abundance.
6. When finished sampling, calculate the percentage of plants infested in the field.

Additional recommendations for monitoring mites in seedling cotton

On seedling cotton (up to 6–8 true leaves) sample regularly to determine the level of infestation using the standard presence/absence technique described above.

When more than 5% of plants are infested it is also advisable to count the numbers of mites on plants, and to score the mite damage level (ie. estimate the % of the plants total leaf area that is damaged by mites).

Continue to monitor mite numbers, damage levels and infestation levels at least weekly, or more frequently if infestation levels are high (> 30% of plants infested).

If the level of infestation, damage level or mite number per plant declines then control is unnecessary, but monitoring should continue.

If mite numbers per plant do not decline after about 6 weeks, if the damage levels exceed an average of 20% of plant leaf area, or if infestation levels increase, then predators are not abundant enough to control mites and a miticide should be applied.

After about 6–8 true leaves, specific mite counts and damage scoring can cease, but continue to use the presence/absence sampling method (points 1–6) until 20% open bolls.

Sampling Tip

...to save time in the field...

Aphids, mites and whitefly can all be sampled using the same leaves from the 3rd or 4th node below the terminal.

Assess for whitefly while collecting the leaves as adults are mobile. Then assess the collected leaves for both mites and aphids.

Collect leaves from several locations in the field.

While the whitefly sampling protocol requires a minimum of 10 leaves per location, aphid and mite sampling requires at least 20 leaves per location. Using 20 leaves will increase the accuracy of whitefly assessment.

Miticide Resistance Monitoring

1. If mites are being collected after a miticide application, ensure sufficient time has lapsed for the miticide to be fully activated. Depending on the product, this may take 7 to 10 days.
2. Collect 50 infested leaves per field. Only collect one sample per field. Keep samples from different fields separate. If mite numbers per leaf are very low, consider collecting up to 100 leaves.
3. Try to avoid collecting all the leaves from only 2 or 3 plants. Where possible collect infested leaves from different areas across the field.
4. Phone Grant Herron and let him know you are sending the sample. Avoid making collections and sending samples on Thursdays or Fridays.
5. Ensure samples are clearly labelled and that labels include the following information:

Farm Name

Field

Region (eg. Gwydir).....

Collector's Name

Phone No

Fax No

Email address.....

Date of collection /..... /.....

Comments eg. details of the problem if a control failure has occurred.

Sending collections to EMAI

Pack the leaves loosely in a paper bag, fold and staple the top. Pack this in a 6-pack esky. Attach the sample details and send by overnight courier to:

Dr Grant Herron
Industry & Investment NSW,
Elizabeth McArthur Agricultural Institute,
Woodbridge Road,
Menangle NSW 2568. Phone: (02) 4640 6471




TABLE 4: Yield reduction caused by mites

The charts below can be used to estimate the percentage of yield reduction caused by mites, for different cotton growing regions.

Days from planting	Current % plants infested with mites																				
	10							30							60						
	Observed rate of increase (%/day)							Observed rate of increase (%/day)							Observed rate of increase (%/day)						
	0.5	1	1.5	2	3	5	7	0.5	1	1.5	2	3	5	7	0.5	1	1.5	2	3	5	7
Warmer regions; planting to 60% bolls open in 134–154 days.																					
Biloela, Bourke, Emerald, Macintyre, Mungindi, St. George, Theodore and Walgett																					
10	1.1	4.0	8.6	14.9	32.8	89.3	100.0	1.8	5.2	17.2	10.3	36.1	94.7	100.0	3.1	7.3	13.2	20.8	41.2	100.0	100.0
20	1.0	3.5	7.4	12.9	28.2	76.7	100.0	1.6	4.6	9.0	14.9	31.2	81.6	100.0	2.6	5.8	10.3	16.0	31.2	76.7	100.0
30	0.9	3.0	6.3	10.9	23.9	65.0	100.0	1.5	4.0	7.8	12.9	26.7	69.6	100.0	2.6	5.8	10.3	16.0	31.2	76.7	100.0
40	0.7	2.5	5.3	9.2	20.0	54.3	100.0	1.3	3.5	6.7	10.9	22.6	58.4	100.0	2.4	5.2	9.0	13.9	26.7	65.0	100.0
50	0.6	2.1	4.4	7.6	16.5	44.5	86.2	1.1	3.0	5.6	9.2	18.8	48.3	91.5	2.2	4.6	7.8	11.9	22.6	54.3	99.6
60	0.5	1.7	3.6	6.1	13.3	35.7	69.1	1.0	2.5	4.7	7.6	15.4	39.1	73.8	2.0	4.0	6.7	10.0	18.8	44.5	81.1
70	0.4	1.4	2.8	4.8	10.4	27.9	53.9	0.9	2.1	3.8	6.1	12.3	30.9	58.0	1.8	3.5	5.6	8.4	15.4	35.7	64.5
80	0.3	1.1	2.2	3.7	7.9	21.0	40.5	0.7	1.7	3.1	4.8	9.5	23.7	44.1	1.6	3.0	4.7	6.8	12.3	27.9	49.9
90	0.3	0.8	1.6	2.7	5.7	15.1	29.1	0.6	1.4	2.4	3.7	7.1	17.4	32.2	1.5	2.5	3.8	5.5	9.5	21.0	37.1
100	0.2	0.6	1.1	1.9	3.9	10.2	19.5	0.5	1.1	2.8	2.7	5.1	12.1	22.1	1.3	2.1	3.1	4.2	7.1	15.1	26.2
110	0.1	0.4	0.7	1.2	2.4	6.3	11.9	0.4	0.8	1.3	1.9	3.4	7.7	13.9	1.1	1.7	2.4	3.2	5.1	10.2	17.2
120	0.1	0.2	0.4	0.6	1.3	3.3	6.1	0.3	0.6	0.8	1.2	2.0	4.3	7.6	1.0	1.4	1.8	2.3	3.4	6.3	10.0
130	0.1	0.1	0.2	0.3	0.5	1.2	2.3	0.3	0.4	0.5	0.6	1.0	1.9	3.2	0.9	1.1	1.3	1.5	2.0	3.3	4.8
140	0.0	0.0	0.0	0.1	0.1	0.2	0.3	0.2	0.2	0.3	0.3	0.3	0.5	0.6	0.7	0.8	0.8	0.9	1.0	1.2	1.5
Average regions; planting to 60% bolls open in 161–170 days.																					
Dalby, Gwydir, Lockyer, Lower Namoi																					
10	1.5	5.3	11.5	20.0	44.1	100.0	100.0	2.3	6.7	13.5	22.6	47.9	100.0	100.0	3.7	9.0	16.7	26.7	53.9	100.0	100.0
20	1.3	4.7	10.1	17.6	38.8	100.0	100.0	2.0	6.0	12.0	20.0	42.3	100.0	100.0	3.4	8.2	15.0	23.9	47.9	100.0	100.0
30	1.2	4.1	8.8	15.4	33.8	92.0	100.0	1.9	5.3	10.6	17.6	37.1	97.4	100.0	3.2	7.4	13.5	21.3	42.3	100.0	100.0
40	1.0	3.6	7.7	13.3	29.1	79.1	100.0	1.7	4.7	9.3	15.4	32.2	84.2	100.0	2.9	6.7	12.0	18.8	37.1	92.0	100.0
50	0.9	3.1	6.5	11.3	24.8	67.3	100.0	1.5	4.1	8.0	13.3	27.6	71.9	100.0	2.7	6.0	10.6	16.5	32.2	79.1	100.0
60	0.8	2.6	5.5	9.5	20.8	56.3	100.0	1.3	3.6	6.9	11.3	23.4	60.6	100.0	2.5	5.3	9.3	14.3	27.6	67.3	100.0
70	0.6	2.2	4.6	7.9	17.2	46.4	89.9	1.2	3.1	5.8	9.5	19.5	50.3	95.2	2.3	4.7	8.0	12.3	23.4	56.3	100.0
80	0.5	1.8	3.7	6.4	13.9	37.4	72.4	1.0	2.6	4.9	7.9	16.0	40.9	77.2	2.0	4.1	6.9	10.4	19.5	46.4	84.7
90	0.4	1.4	3.0	5.1	10.9	29.4	56.8	0.9	2.2	4.0	6.4	12.9	32.5	61.0	1.9	3.6	5.8	8.7	16.0	37.4	67.7
100	0.4	1.1	2.3	3.9	8.4	22.3	43.0	0.8	1.8	3.2	5.1	10.0	25.0	46.8	1.7	3.1	4.9	7.1	12.9	29.4	52.6
110	0.3	0.8	1.7	2.9	6.1	16.2	21.2	0.6	1.4	2.5	3.9	7.6	18.6	34.4	1.5	2.6	4.0	5.7	10.0	22.3	39.5
120	0.2	0.6	1.2	2.0	4.2	11.1	21.3	0.5	1.1	1.9	2.9	5.5	13.1	23.9	1.3	2.2	3.2	4.5	7.6	16.2	28.2
130	0.2	0.4	0.8	1.3	2.7	7.0	13.3	0.4	0.8	1.4	2.0	3.7	8.5	15.4	1.2	1.8	2.5	3.4	5.5	11.1	18.8
140	0.1	0.3	0.5	0.7	1.5	3.8	7.1	0.4	0.6	0.9	1.3	2.3	4.9	8.7	1.0	1.4	1.9	2.4	3.7	7.0	11.3
150	0.1	0.1	0.2	0.3	0.6	1.6	2.9	0.3	0.4	0.6	0.7	1.2	2.3	3.9	0.9	1.1	1.4	1.6	2.3	3.8	5.7
160	0.0	0.0	0.1	0.1	0.2	0.3	0.5	0.2	0.3	0.3	0.3	0.4	0.7	1.0	0.8	0.8	0.9	1.0	1.2	1.6	2.0
Cooler regions; planting to 60% boll open in > 170 days.																					
Boggabri, Breeza, Cecil Plains, Pittsworth, Trangie																					
10	1.7	6.3	13.6	23.7	52.2	100.0	100.0	2.6	7.7	15.7	26.5	56.3	100.0	100.0	4.1	10.2	19.2	30.9	62.8	100.0	100.0
20	1.6	5.6	12.1	21.0	46.4	100.0	100.0	2.3	7.0	14.1	23.7	50.3	100.0	100.0	3.8	9.4	17.4	27.9	56.3	100.0	100.0
30	1.4	4.9	10.7	18.6	40.9	100.0	100.0	2.1	6.3	12.6	21.0	44.5	100.0	100.0	3.5	8.5	15.7	25.0	50.3	100.0	100.0
40	1.2	4.3	9.4	16.2	35.7	97.4	100.0	1.9	5.6	11.1	18.6	39.1	100.0	100.0	3.3	7.7	14.1	22.3	44.5	100.0	100.0
50	1.1	3.8	8.1	14.1	30.9	84.2	100.0	1.7	4.9	9.8	16.2	34.1	89.3	100.0	3.0	7.0	12.6	19.8	39.1	97.4	100.0
60	0.9	3.3	7.0	12.1	26.5	71.9	100.0	1.6	4.3	8.5	14.1	29.4	76.7	100.0	2.8	6.3	11.1	17.4	34.1	84.2	100.0
70	0.8	2.8	5.9	10.2	22.3	60.6	100.0	1.4	3.8	7.3	12.1	25.0	65.0	100.0	2.6	5.6	9.8	15.1	29.4	71.9	100.0
80	0.7	2.3	4.9	8.5	18.6	50.3	97.4	1.2	3.3	6.3	10.2	21.0	54.3	100.0	2.3	4.9	8.5	13.1	25.0	60.6	100.0
90	0.6	1.9	4.1	7.0	15.1	40.9	79.1	1.1	2.8	5.3	8.5	17.4	44.5	84.2	2.1	4.3	7.3	11.1	21.0	50.3	92.0
100	0.5	1.6	3.3	5.6	12.1	32.5	62.8	0.9	2.3	4.3	7.0	14.1	35.7	67.3	1.9	3.8	6.3	9.4	17.4	40.9	74.3
110	0.4	1.2	2.6	4.3	9.4	25.0	48.3	0.8	1.9	3.5	5.6	11.1	27.9	52.2	1.7	3.3	5.3	7.7	14.1	32.5	58.4
120	0.3	0.9	1.9	3.3	7.0	18.6	35.7	0.7	1.6	2.8	4.3	8.5	21.0	39.1	1.5	2.8	4.3	6.3	11.1	25.0	44.5
130	0.2	0.7	1.4	2.3	4.9	13.1	25.0	0.6	1.2	2.1	3.3	6.3	15.1	27.9	1.4	2.3	3.5	4.9	8.5	18.6	32.5
140	0.2	0.5	0.9	1.6	3.3	8.5	16.2	0.5	0.9	1.6	2.3	4.3	10.2	18.6	1.2	1.9	2.8	3.8	6.3	13.1	22.3
150	0.1	0.3	0.6	0.9	1.9	4.9	9.4	0.4	0.7	1.1	1.6	2.8	6.3	11.1	1.1	1.6	2.1	2.8	4.3	8.5	14.1
160	0.1	0.2	0.3	0.5	0.9	2.3	4.3	0.3	0.5	0.7	0.9	1.6	3.3	5.6	0.9	1.2	1.6	1.9	2.8	4.9	7.7
170	0.0	0.1	0.1	0.2	0.3	0.7	1.2	0.2	0.3	0.4	0.5	0.7	1.2	1.9	0.8	0.9	1.1	1.2	1.6	2.3	3.3

MITE YIELD LOSS ESTIMATOR CHARTS






TABLE 5: Control of mites

Active ingredient	Concentration and formulation	Application rate of product	Comments
Mite (<i>Tetranychus</i>) species			
Abamectin	18 g/L EC	0.3 L/ha	Best results will be obtained when applied to low mite populations. Do not use more than twice in one season.
Aldicarb	150 g/kg G	3.0–7.0 kg/ha	Apply into the seed furrow at sowing.
Amitraz	200 g/L EC	2.0 L/ha	Suppression when used for controlling <i>Helicoverpa</i>
Bifenthrin	100 g/L EC 250g/L EC	0.6–0.8 L/ha 0.24-0.32L/ha	Applications against <i>Helicoverpa</i> spp. will give good control of low mite populations.
Chlorpyrifos	300 g/L EC	1.0–1.5 L/ha 2.5 L/ha	Mix with pyrethroids as a preventative spray to minimise buildup of mite populations. For established mite populations.
Diafenthion	500 g/L SC	0.6 or 0.8 L/ha	Treatment at higher infestation levels may lead to unsatisfactory results.
Dicofol	240 g/L EC	4.0 L/ha	NSW registration only. Apply by ground rig at first appearance of mites before row closure.
Dimethoate	400 g/L EC	0.34–0.5 L/ha	Will not control organophosphate-resistant mites.
Disulfoton	50 g/kg G	14.0 kg/ha	Apply in a band near seed at planting.
Emamectin benzoate	17 g/L EC	0.55–0.7 L/ha	When applied for <i>Helicoverpa</i> control will reduce the rate of mite population development. Suppression only.
Etoxazole	110 g/L SC	0.35 L/ha	Apply by ground rig only. Best results when used on low to increasing populations.
Methidathion	400 g/L EC	1.4 L/ha	Knockdown and short residual control.
Phorate	100 g/kg G	6.0 kg/ha 11.0–17.0 kg/ha	For short residual control. For extended period of control. Only use the highest rate on heavy soils when conditions favour good emergence.
	200 g/kg G	3.0 kg/ha 5.5–8.5 kg/ha	For short residual control. NSW registration only.
Profenofos	250 g/L EC 500 g/L EC	1.0–2.0 L/ha 0.5–1.0 L/ha	Use low rate to suppress build up. Use high rate if aphids present.
Propargite	600 g/L EC	2.5 L/ha	Apply as spray before mite infestations reach damaging levels.